

## WEST Search History





DATE: Monday, May 01, 2006

Hide?	Set Name	Query	Hit Count
<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L19	L18 and l17	14
<input type="checkbox"/>	L18	l16 and selenium	32
<input type="checkbox"/>	L17	L16 and retinoid	21
<input type="checkbox"/>	L16	L15 and treatment	84
<input type="checkbox"/>	L15	L14 and glutathione adj peroxidase	86
<input type="checkbox"/>	L14	HCV	7569
<input type="checkbox"/>	L13	Munchen S.in.	1
<i>DB=EPAB; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L12	DE-10255861-A1.did.	1
<input type="checkbox"/>	L11	DE-10255861-A1.did.	1
<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L10	L9 and glutathione adj peroxidase	3
<input type="checkbox"/>	L9	Cotten m.in.	32
<input type="checkbox"/>	L8	L7 and glutathione adj peroxidase	5
<input type="checkbox"/>	L7	Herget T.in.	10
<input type="checkbox"/>	L6	L5 and glutathione adj peroxidase	1
<input type="checkbox"/>	L5	L1 and HCV	61
<input type="checkbox"/>	L4	L1 and l2	5
<input type="checkbox"/>	L3	424/228.1.ICLS.	133
<input type="checkbox"/>	L2	424/228.1.ICLS.	133
<input type="checkbox"/>	L1	424/93.2.ICLS.	1637

END OF SEARCH HISTORY

## WEST Search History





DATE: Monday, May 01, 2006

Hide?	Set Name	Query	Hit Count
<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L19	L18 and l17	14
<input type="checkbox"/>	L18	l16 and selenium	32
<input type="checkbox"/>	L17	L16 and retinoid	21
<input type="checkbox"/>	L16	L15 and treatment	84
<input type="checkbox"/>	L15	L14 and glutathione adj peroxidase	86
<input type="checkbox"/>	L14	HCV	7569
<input type="checkbox"/>	L13	Munchen S.in.	1
<i>DB=EPAB; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L12	DE-10255861-A1.did.	1
<input type="checkbox"/>	L11	DE-10255861-A1.did.	1
<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L10	L9 and glutathione adj peroxidase	3
<input type="checkbox"/>	L9	Cotten m.in.	32
<input type="checkbox"/>	L8	L7 and glutathione adj peroxidase	5
<input type="checkbox"/>	L7	Herget T.in.	10
<input type="checkbox"/>	L6	L5 and glutathione adj peroxidase	1
<input type="checkbox"/>	L5	L1 and HCV	61
<input type="checkbox"/>	L4	L1 and l2	5
<input type="checkbox"/>	L3	424/228.1.ICLS.	133
<input type="checkbox"/>	L2	424/228.1.ICLS.	133
<input type="checkbox"/>	L1	424/93.2.ICLS.	1637

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1648BQL

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/  
USPAT2  
NEWS 4 JAN 13 IPC 8 searching in IFIPAT, IFIUDB, and IFICDB  
NEWS 5 JAN 13 New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to  
INPADOC  
NEWS 6 JAN 17 Pre-1988 INPI data added to MARPAT  
NEWS 7 JAN 17 IPC 8 in the WPI family of databases including WPIFV  
NEWS 8 JAN 30 Saved answer limit increased  
NEWS 9 FEB 21 STN AnaVist, Version 1.1, lets you share your STN AnaVist  
visualization results  
NEWS 10 FEB 22 The IPC thesaurus added to additional patent databases on STN  
NEWS 11 FEB 22 Updates in EPFULL; IPC 8 enhancements added  
NEWS 12 FEB 27 New STN AnaVist pricing effective March 1, 2006  
NEWS 13 FEB 28 MEDLINE/LMEDLINE reload improves functionality  
NEWS 14 FEB 28 TOXCENTER reloaded with enhancements  
NEWS 15 FEB 28 REGISTRY/ZREGISTRY enhanced with more experimental spectral  
property data  
NEWS 16 MAR 01 INSPEC reloaded and enhanced  
NEWS 17 MAR 03 Updates in PATDPA; addition of IPC 8 data without attributes  
NEWS 18 MAR 08 X.25 communication option no longer available after June 2006  
NEWS 19 MAR 22 EMBASE is now updated on a daily basis  
NEWS 20 APR 03 New IPC 8 fields and IPC thesaurus added to PATDPAFULL  
NEWS 21 APR 03 Bibliographic data updates resume; new IPC 8 fields and IPC  
thesaurus added in PCTFULL  
NEWS 22 APR 04 STN AnaVist \$500 visualization usage credit offered  
NEWS 23 APR 12 LINSPEC, learning database for INSPEC, reloaded and enhanced  
NEWS 24 APR 12 Improved structure highlighting in FQHIT and QHIT display  
in MARPAT  
NEWS 25 APR 12 Derwent World Patents Index to be reloaded and enhanced during  
second quarter; strategies may be affected  
  
NEWS EXPRESS FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a,  
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.  
V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT  
<http://download.cas.org/express/v8.0-Discover/>  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items  
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \*

COMPLETE THE STN SURVEY - APRIL 27 THROUGH MAY 31

Dear valued STN customer,

In an effort to enhance your experience with STN, we would like to better understand what you find useful. Please take approximately 5 minutes to complete a web survey.

If you provide us with your name, login ID, and e-mail address, you will be entered in a drawing to win a free iPod(R). Your responses will be kept confidential and will help us make future improvements to STN.

Take survey: <http://www.zoomerang.com/survey.zgi?p=WEB2259HNKWTUW>

Thank you in advance for your participation.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 09:17:47 ON 01 MAY 2006

=> file caplus biosis  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 09:18:11 ON 01 MAY 2006  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 09:18:11 ON 01 MAY 2006  
Copyright (c) 2006 The Thomson Corporation

=> "glutathione peroxidase"  
L1 30666 "GLUTATHIONE PEROXIDASE"

=> HCV  
L2 33362 HCV

=> L1 and L2  
L3 28 L1 AND L2

=> gastroiintestinal  
L4 0 GASTROIINTESTINAL

=> gastrointestinal  
L5 184440 GASTROINTESTINAL

=> L5 and L3  
L6 9 L5 AND L3

=> D L6 IBIB ABS 1-9

L6 ANSWER 1 OF 9 CAPLUS COPYRIGHT 2006 ACS on STN  
ACCESSION NUMBER: 2005:1331259 CAPLUS  
DOCUMENT NUMBER: 144:64327  
TITLE: Use of selenium or a selenium salt and a retinoid acid  
or a retinoid in the treatment of viral hepatitis C  
INVENTOR(S): Herget, Thomas; Klebl, Bert  
PATENT ASSIGNEE(S): GPC Biotech A.-G., Germany  
SOURCE: PCT Int. Appl., 58 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005120479	A1	20051222	WO 2005-EP6226	20050609
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GR, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: US 2004-578161P P 20040609

AB The present invention relates to combination therapies comprising at least one retinoid or retinoid agonist together with selenium or a selenium salt particularly useful in conjunction with conventional antiviral therapeutics which are synergistically effective against Hepatitis C virus (HCV) infections. In particular, the present invention relates to the synergism between compds. capable of activating or upregulating the **gastrointestinal** form of **glutathione peroxidase** for prophylaxis and/or treatment of HCV infections, administered in combination therapies with interferons. The combinations disclosed have proven surprisingly effective even in patients unresponsive to interferon/ribavirin therapies.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:204131 CAPLUS

DOCUMENT NUMBER: 142:277684

TITLE: Expression of **Gastrointestinal Glutathione Peroxidase** Is Inversely Correlated to the Presence of Hepatitis C Virus Subgenomic RNA in Human Liver Cells

AUTHOR(S): Morbitzer, Monika; Herget, Thomas  
 CORPORATE SOURCE: AXXIMA Pharmaceuticals AG, Munich, 81377, Germany  
 SOURCE: Journal of Biological Chemistry (2005), 280(10), 8831-8841

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular Biology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB There is great medical need to develop novel therapies for treatment of human hepatitis C virus (HCV). By gene expression anal. of three HCV-subgenomic RNA replicon cell lines, we identified cellular proteins whose expression is affected by the presence of HCV and therefore may serve as drug targets. Data from cDNA array filter hybridization, as well as from Northern and Western blotting, revealed that the **gastrointestinal-glutathione peroxidase** (GI-GPx) was drastically down-regulated (up to 20-fold) in all replicon cell lines tested. Concomitantly, total cellular **glutathione peroxidase** activity was drastically reduced, which rendered these human liver cells more susceptible toward oxidative stress. Interferon  $\alpha$  caused down-regulation of the HCV-replicon followed by recovery of GI-GPx expression to nearly normal levels. Furthermore, expression of GI-GPx in replicon cells by gene transduction caused down-regulation of HCV RNA in a dose-dependent manner. Moreover, activating the endogenous gene coding for GI-GPx by all-trans-retinoic acid (RA) was sufficient to cause down-regulation of the HCV replicon. A small interfering RNA duplex abrogated GI-GPx up-regulation by RA and concomitantly suppression of HCV. The RA effect was dependent on the presence of sodium selenite, was reversible, and was independent of RNA-activated protein kinase. Taken together, these results show that HCV inhibits the expression of GI-GPx in replicon cells to promote its intracellular

propagation. Modulation of GI-GPx activity may open new avenues of treatment for HCV patients.

REFERENCE COUNT: 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 3 OF 9 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:633154 CAPLUS

DOCUMENT NUMBER: 141:167729

TITLE: **Gastrointestinal glutathione peroxidase** as therapeutic target for treatment of HCV infection, methods of treating HCV infection, and compounds useful therefor

INVENTOR(S): Herget, Thomas; Cotten, Matthew; Obert, Sabine; Klebl, Bert

PATENT ASSIGNEE(S): Germany

SOURCE: U.S. Pat. Appl. Publ., 24 pp., Cont.-in-part of U.S. Pat. Appl. 2003 180,719.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004152073	A1	20040805	US 2003-723719	20031126
WO 2002084294	A2	20021024	WO 2002-EP4167	20020415
WO 2002084294	A3	20031030		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
DE 10255861	A1	20040617	DE 2002-10255861	20021129
US 2003180719	A1	20030925	US 2003-342054	20030114
PRIORITY APPLN. INFO.:			US 2001-283345P	P 20010413
			WO 2002-EP4167	A2 20020415
			DE 2002-10255861	A 20021129
			US 2002-430367P	P 20021203
			US 2003-342054	A2 20030114

AB The present invention relates to the human cellular protein **glutathione peroxidase-gastrointestinal** as a target for medical intervention against Hepatitis C virus (HCV) infections. Furthermore, the present invention relates to a method for the detection of compds. useful for prophylaxis and/or treatment of hepatitis C virus infections and a method for detecting hepatitis C virus infections in an individual or in cells. Also compns., compds., nucleic acid mols. (such as aptamers), mono- or polyclonal antibodies are disclosed which are effective for the treatment of HCV infections, and methods for prophylaxis and/or treatment of hepatitis C virus infections or for the regulation of hepatitis C virus production are disclosed. The inventors designed a randomized, single-blinded clin. study to test the safety, tolerability, and efficacy of all-trans retinoic acid alone or in combination with pegylated  $\alpha$  interferon in patients with chronic hepatitis C. The therapy regimens include: Vesanoid (orally administered all-trans retinoic acid compound, Hoffman-La Roche); Pegasys (slow-release pegylated interferon  $\alpha$ 2a, Hoffman-La Roche); and selen 30 ALLACT (supplement containing selenium and ALLACT composed of garlic powder and Lactobacillus bulgaricus).

L6 ANSWER 4 OF 9 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:490732 CAPLUS

DOCUMENT NUMBER: 141:42933

TITLE: Formulations useful against hepatitis C virus

infections

INVENTOR(S): Herget, Thomas; Klebl, Bert  
PATENT ASSIGNEE(S): Axxima Pharmaceuticals A.-G., Germany  
SOURCE: PCT Int. Appl., 72 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 4  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004050101	A2	20040617	WO 2003-EP13514	20031201
WO 2004050101	A3	20040910		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
DE 10255861	A1	20040617	DE 2002-10255861	20021129
DE 10305138	A1	20040826	DE 2003-10305138	20030207
CA 2509955	AA	20040617	CA 2003-2509955	20031201
AU 2003294757	A1	20040623	AU 2003-294757	20031201
EP 1567172	A2	20050831	EP 2003-785699	20031201
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
JP 2006514094	T2	20060427	JP 2004-570683	20031201
PRIORITY APPLN. INFO.:			DE 2002-10255861	A 20021129
			US 2002-430367P	P 20021203
			DE 2003-10305138	A 20030207
			US 2003-446246P	P 20030211
			WO 2003-EP13514	W 20031201

AB The present invention relates generally to chemical compds. and substances which are effective against Hepatitis C virus (HCV) infections. Moreover, the present invention relates to compns. comprising said compds. and/or substances, to methods for preventing HCV infections as well use of the compds. and/or substances for the preparation of compns. useful for the prophylaxis and/or treatment of HCV infections. Useful compds. and substances according to the invention are selenium, selenium salts, Vitamin D3 and retinoids, like all trans retinoic acid and salts thereof, C1-C10 alkyl amide of all trans retinoic acid and salts thereof, C1-C10 alkyl esters of all trans retinoic acid and salts thereof, 9-cis retinoic acid and salts thereof, C1-C10 alkyl amide of 9-cis retinoic acid and salts thereof, C1-C10 alkyl esters of 9-cis retinoic acid and salts thereof, (E)-4-[2-(5,6,7,8-tetrahydro-5,5,8,8-tetra methyl-2-naphthalenyl-1)-propenyl] benzoic acid (TTNPB), (4-[5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl] carboxamido) benzoic acid (AM-580), N-(4-hydroxyphenyl) retinamide (4-HPR), and 6-[3-(1-adamantyl)-4-hydroxyphenyl]-2-naphthalene carboxylic acid (AHPN).

L6 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:757185 CAPLUS  
DOCUMENT NUMBER: 139:271014  
TITLE: Human cellular protein **gastrointestinal glutathione peroxidase** as target for medical intervention against hepatitis C virus infections  
INVENTOR(S): Herget, Thomas; Cotten, Matthew; Obert, Sabine  
PATENT ASSIGNEE(S): Germany  
SOURCE: U.S. Pat. Appl. Publ., 23 pp., Cont.-in-part of Appl. No. PCT/EP02/04167.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2003180719	A1	20030925	US 2003-342054	20030114
WO 2002084294	A2	20021024	WO 2002-EP4167	20020415
WO 2002084294	A3	20031030		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
DE 10255861	A1	20040617	DE 2002-10255861	20021129
US 2004152073	A1	20040805	US 2003-723719	20031126
PRIORITY APPLN. INFO.:			US 2001-283345P	P 20010413
			WO 2002-EP4167	A2 20020415
			DE 2002-10255861	A 20021129
			US 2002-430367P	P 20021203
			US 2003-342054	A2 20030114

AB The present invention relates to the human cellular protein **glutathione peroxidase-gastrointestinal** as a target for medical intervention against Hepatitis C virus (HCV) infections. Furthermore, the present invention relates to a method for the detection of compds. useful for prophylaxis and/or treatment of Hepatitis C virus infections and a method for detecting Hepatitis C virus infections in an individual or in cells. Also compns., compds., nucleic acid mols. (such as aptamers), mono- or polyclonal antibodies are disclosed which are effective for the treatment of **HCV** infections, and methods for prophylaxis and/or treatment of Hepatitis C virus infections or for the regulation of Hepatitis C virus production are disclosed.

L6 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:814448 CAPLUS

DOCUMENT NUMBER: 137:291285

TITLE: Human cellular protein **gastrointestinal glutathione peroxidase** as target for medical intervention against hepatitis c virus infections

INVENTOR(S): Herget, Thomas; Cotten, Matthew; Obert, Sabine

PATENT ASSIGNEE(S): Axxima Pharmaceuticals Ag, Germany

SOURCE: PCT Int. Appl., 37 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002084294	A2	20021024	WO 2002-EP4167	20020415
WO 2002084294	A3	20031030		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2443525	AA	20021024	CA 2002-2443525	20020415



EP 1377833	A2	20040107	EP 2002-730159	20020415
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004533822	T2	20041111	JP 2002-581997	20020415
US 2003180719	A1	20030925	US 2003-342054	20030114
US 2004152073	A1	20040805	US 2003-723719	20031126
PRIORITY APPLN. INFO.:			US 2001-283345P	P 20010413
			WO 2002-EP4167	W 20020415
			DE 2002-10255861	A 20021129
			US 2002-430367P	P 20021203
			US 2003-342054	A2 20030114

AB The present invention relates to the human cellular protein **glutathione peroxidase-gastrointestinal** as potential targets for medical intervention against Hepatitis C virus (HCV) infections. Furthermore, the present invention relates to a method for the detection of compds. useful for prophylaxis and/or treatment of Hepatitis C virus infections and a method for detecting Hepatitis C virus infections in an individual or in cells. Also mono- or polyclonal antibodies are disclosed effective for the treatment of HCV infections together with methods for treating Hepatitis C virus infections or for the regulation of Hepatitis C virus production wherein said antibodies may be used.

L6 ANSWER 7 OF 9 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN  
 ACCESSION NUMBER: 2006:210744 BIOSIS  
 DOCUMENT NUMBER: PREV200600212473  
 TITLE: Retinoic acid causes up-regulation of the **gastrointestinal glutathione peroxidase** (GI-GPx) promoter and concomitantly down-regulation of hepatitis C virus (HCV) subgenomic RNA.  
 AUTHOR(S): Herget, T.; Morbitzer, M.; Klebl, B.; Galle, Peter; Becher, Wulf; Wallasch, Christian  
 SOURCE: Gastroenterology, (APR 2005) Vol. 128, No. 4, Suppl. 2, pp. A699.  
 Meeting Info.: Annual Meeting of the American-Gastroenterological-Association/Digestive-Disease-Week. Chicago, IL, USA. May 14 -19, 2005. Amer Gastroenterol Assoc.  
 CODEN: GASTAB. ISSN: 0016-5085.  
 DOCUMENT TYPE: Conference; (Meeting)  
 Conference; Abstract; (Meeting Abstract)  
 LANGUAGE: English  
 ENTRY DATE: Entered STN: 29 Mar 2006  
 Last Updated on STN: 29 Mar 2006

AB The mRNA expression patterns of three Hepatitis C Virus (HCV) )-subgenomic RNA replicon cell lines were compared with those of mock transfected or untransfected HuH7 cells utilizing cDNA array filters. The **gastrointestinal-glutathione peroxidase** (GI-GPx) mRNA was drastically down-regulated (as low as 5 to 10% of controls) in all replicon cell lines, while the expression level of the classical cellular-**glutathione peroxidase** (cGPx) remained unaffected. These data were confirmed by Northern blot and Western blot analyses. GI-GPx is a selenoprotein belonging to a family of four members, responsible for the detoxification of peroxides. Measuring total cellular **glutathione peroxidase** activity, revealed that the replicon cells showed reduced **glutathione peroxidase** activity (approx. 50% of control cells). Accordingly, replicon cells demonstrated increased susceptibility towards paraquat, a compound producing oxidative stress, reflected by a reduced viability of the replicon cultures compared to mock-transfected cell lines. When replicon cells were incubated with interferon for four days to induce the innate immune response, the HCV-replicon became down-regulated. Concomitantly, expression of CI-GPx resumed to nearly normal levels. Interferon itself did not effect the expression of GI-GPx in mock transfected and naive HuH7 cells. Furthermore, transient over-expression of the GI-GPx cDNA via adenoviral gene transfer induced a substantial and consistent down-regulation of the HCV RNA and the NS5a protein in replicon cells. In depth inspection of the 5' promoter region of the

GI-GPx gene revealed the presence of two retinoic acid response elements (RARE). Treating replicon cultures with retinoic acid in the presence of selenite lead to increased expression of endogenous GI-GPx, followed by a dramatic down-regulation of the replicon. This decrease was even more pronounced, when cells were incubated with retinoic acid in the presence of selenite and interferon alpha. Taken together, these data show, that (a) expression of GI-GPx and replication of **HCV** exclude each other and (b) retinoic acid might be a valuable tool for the treatment of **HCV** patients. Therefore, a clinical pilot trial at the University of Mainz with 9 population of interferon non-responders was initiated. Preliminary data of this clinical trial will be presented in parallel.

L6 ANSWER 8 OF 9 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN  
ACCESSION NUMBER: 2006:210736 BIOSIS  
DOCUMENT NUMBER: PREV200600212465  
TITLE: All-trans-retinoic acid for treatment of patients with chronic hepatitis C and non-response to interferon alfa/ribavirin.  
AUTHOR(S): Becher, Wulf O.; Wallasch, Christian; Herget, T.; Klebl, B. M.; Galle, Peter R.; Strand, D.  
SOURCE: Gastroenterology, (APR 2005) Vol. 128, No. 4, Suppl. 2, pp. A697-A698.  
Meeting Info.: Annual Meeting of the American-Gastroenterological-Association/Digestive-Disease-Week. Chicago, IL, USA. May 14 -19, 2005. Amer Gastroenterol Assoc.  
CODEN: GASTAB. ISSN: 0016-5085.  
DOCUMENT TYPE: Conference; (Meeting)  
Conference; Abstract; (Meeting Abstract)  
LANGUAGE: English  
ENTRY DATE: Entered STN: 29 Mar 2006  
Last Updated on STN: 29 Mar 2006

AB Introduction: In vitro studies, submitted in parallel by Herget et al, have shown that all-trans retinoic acid (ATRA) induces upregulation of selenium dependent **gastrointestinal-glutathione peroxidase** in **HCV**-subgenomic RNA replicon cells leading to drastic downregulation of the replicon, that was further enhanced by interferon alfa. Based on these findings, a clinical pilot trial was performed in **HCV** non-responder patients. Methods: 20 patients with chronic **HCV** infection and non-response to IFN alfa and ribavirin (pos. PCR at week 12) were randomly assigned to treatment with daily 45 mg/m<sup>2</sup> ATRA p.o. and 30 mcg/d selenite (arm A) or 45 mg/m<sup>2</sup> ATRA and selenite combined with 180 mcg/week peg-interferon alfa2a (arm B). All patients had serotype-1, elevated ALT levels and 9 patients had F3 fibrosis or cirrhosis. Mean IFNa pretreatment duration was 14 months, 9 patients were Peg-IFN nonresponders. ATRA treatment was continued for 12 weeks and followed for additional 12 weeks after end of treatment (ETR). **HCV** RNA was assessed by quantitative real time PCR.

L6 ANSWER 9 OF 9 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN  
ACCESSION NUMBER: 2005:213630 BIOSIS  
DOCUMENT NUMBER: PREV200510004953  
TITLE: Expression of **gastrointestinal glutathione peroxidase** is inversely correlated to the presence of hepatitis C virus subgenomic RNA in human liver cells.  
AUTHOR(S): Morbitzer, Monika; Herget, Thomas [Reprint Author]  
CORPORATE SOURCE: Merck KGaA, Frankfurter Str 250, D-64293 Darmstadt, Germany  
thomas.herget@merck.de  
SOURCE: Journal of Biological Chemistry, (MAR 11 2005) Vol. 280, No. 10, pp. 8831-8841.  
CODEN: JBCHA3. ISSN: 0021-9258.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 10 Jun 2005  
Last Updated on STN: 10 Jun 2005

AB There is great medical need to develop novel therapies for treatment of human hepatitis C virus (**HCV**). By gene expression analysis of three **HCV**-subgenomic RNA replicon cell lines, we identified

cellular proteins whose expression is affected by the presence of **HCV** and therefore may serve as drug targets. Data from cDNA array filter hybridization, as well as from Northern and Western blotting, revealed that the **gastrointestinal-glutathione peroxidase** (GI-GPx) was drastically down-regulated (up to 20-fold) in all replicon cell lines tested. Concomitantly, total cellular **glutathione peroxidase** activity was drastically reduced, which rendered these human liver cells more susceptible toward oxidative stress. Interferon alpha caused down-regulation of the **HCV** -replicon followed by recovery of GI-GPx expression to nearly normal levels. Furthermore, expression of GI-GPx in replicon cells by gene transduction caused down-regulation of **HCV** RNA in a dose-dependent manner. Moreover, activating the endogenous gene coding for GI-GPx by all-trans-retinoic acid (RA) was sufficient to cause down-regulation of the **HCV** replicon. A small interfering RNA duplex abrogated GI-GPx up-regulation by RA and concomitantly suppression of **HCV**. The RA effect was dependent on the presence of sodium selenite, was reversible, and was independent of RNA-activated protein kinase. Taken together, these results show that **HCV** inhibits the expression of GI-GPx in replicon cells to promote its intracellular propagation. Modulation of GI-GPx activity may open new avenues of treatment for **HCV** patients.